



VETERINARY PRODUCTS LABORATORY

IMPORT - EXPORT

Address: Av. B. Ader 3205 - (1605) Munro - Buenos Aires - República Argentina

Phone/Fax: 0054 11-4762-5163

E-mail: info@chinfield.com

Visit our Web site: www.chinfield.com

TECHNICAL DATA SHEET

Product Name: FUROFIELD®

FORMULA:

Furosemide	5.000 g
Dexamethasone 21 Phosphate	0.200 g
Formulation agents q.s. to	100 ml

TARGET SPECIES:

Equines. Cattle. Dogs. Cats.

INDICATIONS:

Bronchospasm. Bronchitis. Bronchiolitis. Chronic Obstructive Pulmonary Disease (COPD). Respiratory allergies of different origins.

As an adjuvant in the antibiotic treatment of respiratory infections.

Its activity is effective both in acute and chronic respiratory diseases, allowing its use in symptomatic emergency therapy or in long-term treatments.

PRESENTATION:

Amber glass vials of 25 ml.

APPEARANCE:

Clear, slightly yellowish liquid solution.

DOSAGE AND ADMINISTRATION:

EQUINES AND CATTLE: 10 to 25 ml depending on the professional's judgment and the problem to be treated.

DOGS AND CATS: 0.25 to 0.5 ml depending on the professional's judgment and the problem to be treated.

Routes of administration: Intramuscular. Intravenous injection. Intravenous injection. Oral route only for equines and dogs.

CONTRAINDICATIONS:

Hypersensitivity. Acute systemic infections (unless specific antimicrobial therapy is established). Avoid live attenuated vaccines. Should not be used in patients with active or possibly inactive tuberculosis except when used in conjunction with tuberculostatic drugs. Acute glomerulonephritis or acute renal failure.



VETERINARY PRODUCTS LABORATORY

IMPORT - EXPORT

Address: Av. B. Ader 3205 - (1605) Munro - Buenos Aires - República Argentina

Phone/Fax: 0054 11-4762-5163

E-mail: info@chinfield.com

Visit our Web site: www.chinfield.com

SIDE EFFECTS:

Interactions:

- Amphotericin B or potassium-depleting diuretics: concomitant use may cause hypokalemia.
- Digitalis glycosides: may increase the likelihood of digitalis intoxication associated with hypokalemia.
- Acetylsalicylic acid: should be used with caution, especially in states of hypoprothrombinemia.
- Isoniazid: its antituberculous effectiveness may be reduced due to increased hepatic metabolism, excretion of isoniazid, or both, leading to decreased plasma concentration, especially in patients who are rapid acetylators.

WARNINGS:

Keep out of reach of children and pets. National Poison Control Center: 0800-333-0160 Buenos Aires, Argentina.

Must be administered by a veterinarian.

Chinfield S.A. is not responsible for problems caused by improper use and/or storage of the product.

STORAGE CONDITIONS:

Store below 30 °C, protected from light.

Do not refrigerate. Do not freeze.

SALES UNDER PRESCRIPTION ONLY

Manufactured and Distributed by: Chinfield S.A. Av. Ader 3205, Munro, Buenos Aires, Argentina

E-Mail: asuntosregulatorios@chinfield.com

www.chinfield.com