



VETERINARY PRODUCTS LABORATORY

IMPORT - EXPORT

Address: Av. B. Ader 3205 - (1605) Munro - Buenos Aires - República Argentina

Phone/Fax: 0054 11-4762-5163

E-mail: info@chinfield.com

Visit our Web site: www.chinfield.com

TECHNICAL DATA SHEET

Product Name: TRIAMCINOLONA CHINFIELD®

FORMULA:

Triamcinolone Acetonide 0.600 g
Formulation agents q.s. to 100 ml

TARGET SPECIES:

Equines, cattle, canines, felines.

INDICATIONS:

Indicated in pruritus, eczema, allergic processes, cellulitis, arthritis, synovitis, bursitis, tenosynovitis, rheumatoid syndrome, blepharitis, bovine ketosis, hepatitis, lymphosarcoma, aseptic inflammatory processes, traumatic inflammation.

PRESENTATION:

Printed cardboard box containing a vial, with label and leaflet.
Content: 5 ml.

APPEARANCE:

Viscous white suspension.

DOSAGE AND ADMINISTRATION:

Cattle, Canines, and Felines: Subcutaneous, Intramuscular, or Intrasynovial.
Equines: Intramuscular or Intrasynovial (NOT SUBCUTANEOUS).

Doses:

Equines: 0.6 ml per 100 kg b.w.

Cattle: 0.6 ml per 100 kg b.w.

Canines: 0.4 ml per 10 kg b.w.

Felines: 0.3 ml total.

CONTRAINDICATIONS:

In cases of hypersensitivity. Do not administer intravenously. Do not use in septic processes unless appropriate antimicrobial therapy is established. Do not use in fungal infections. Should not be used in patients with active or suspected inactive tuberculosis except when used with tuberculostatic drugs. Do not use when vaccines of attenuated organisms are applied. Contraindicated in amyloidosis, mange, hepatic insufficiency, renal insufficiency, ulcerative colitis. Do not apply during healing of major wounds as this drug delays tissue repair.



VETERINARY PRODUCTS LABORATORY

IMPORT - EXPORT

Address: Av. B. Ader 3205 - (1605) Munro - Buenos Aires - República Argentina

Phone/Fax: 0054 11-4762-5163

E-mail: info@chinfield.com

Visit our Web site: www.chinfield.com

Do not use in females during the last third of gestation due to the possibility of premature birth. In equines: contraindicated in gastric and/or duodenal ulcers. Do not use in laminitis. Do not apply intra-articularly in septic joint processes or articular fractures. Contraindicated in diabetic animals.

SIDE EFFECTS:

Hypercorticism caused by continuous high doses of corticosteroids for a prolonged period.

Hypocorticism due to sudden withdrawal of the drug after prolonged treatment.

Delayed wound healing processes.

Reproduction: do not use in pregnant females in the last third of gestation due to risk of premature birth.

High doses and long treatments may cause or aggravate laminitis.

High doses and long treatments may increase hydrochloric acid and pepsin production in the stomach, resulting in gastric ulcers.

WARNINGS:

Keep out of reach of children and pets. National Poison Control Center: 0800-333-0160 Buenos Aires, Argentina.

Must be administered by a veterinarian.

Chinfield S.A. is not responsible for problems caused by improper use and/or storage of the product.

STORAGE CONDITIONS:

Store below 30 °C, protected from light.

Do not refrigerate. Do not freeze.

SALES UNDER PRESCRIPTION ONLY

Manufactured and Distributed by: Chinfield S.A. Av. Ader 3205, Munro, Buenos Aires, Argentina

E-Mail: asuntosregulatorios@chinfield.com

www.chinfield.com