



VETERINARY PRODUCTS LABORATORY

IMPORT - EXPORT

Address: Av. B. Ader 3205 - (1605) Munro - Buenos Aires - República Argentina

Phone/Fax: 0054 11-4762-5163

E-mail: info@chinfield.com

Visit our Web site: www.chinfield.com

TECHNICAL DATA SHEET

Product Name: NABOLIC STRONG®

FORMULA:

Stanozolol 5.000 g

Formulation agents q.s. to 100 ml

TARGET SPECIES:

Equines.

INDICATIONS:

Indicated for the recovery of organisms weakened by advanced age, diseases, and their convalescence, traumatic injuries or excessive work. Useful in cases of fractures, orthopedic conditions, muscular dystrophies, and general bone deficiencies. Also indicated as an appetite stimulant and to increase strength and vitality.

PRESENTATION:

The product is presented in 30 ml colorless glass vials with label, box, and leaflet.

APPEARANCE:

Viscous white suspension.

DOSAGE AND ADMINISTRATION:

1 or 2 ml per 100 kg b.w., according to the application interval:

- 1 dose per week: 1 ml per 100 kg b.w.
- 1 dose every 15 days: 2 ml per 100 kg b.w.

Route of administration: Intramuscular.

CONTRAINDICATIONS:

Do not overdose.

Do not administer to pregnant mares.

Do not administer to stallions during breeding season.

Do not administer to horses intended for human consumption.

Do not administer to growing animals, as it may stop growth by accelerating epiphyseal closure. In cases of malnutrition, convalescence from infectious diseases, growth disorders, etc., consult the veterinarian. In such cases, periodic radiographic controls (every 30 days) of growth plates are recommended to verify that premature ossification does not occur. This contraindication does not apply to horses over 24 months of age.



VETERINARY PRODUCTS LABORATORY

IMPORT - EXPORT

Address: Av. B. Ader 3205 - (1605) Munro - Buenos Aires - República Argentina

Phone/Fax: 0054 11-4762-5163

E-mail: info@chinfield.com

Visit our Web site: www.chinfield.com

Do not administer in cases of malignant tumors of any etiology.

Animals with renal and/or cardiac deficiencies should be monitored by the veterinarian during treatment, as anabolic therapy may be accompanied by sodium and water retention.

SIDE EFFECTS:

No side effects or incompatibilities when applied and dosed as indicated.

At the recommended doses, no undesirable side effects were observed.

These side effects may appear in cases of overdose and prolonged treatments, showing signs of virilization, cholestatic hepatotoxicity clinically manifested by anorexia, skin disorders, constipation, halitosis, flatulence, tympanism, etc., and biochemically characterized by an increase in alkaline phosphatase, bilirubin, total cholesterol, and GGT. Sodium and water retention with edema production, as well as diffuse alopecia localized on the rib cage, are also possible. These processes are reversible once the drug is discontinued.

In terms of incompatibilities, adrenocorticotrophic hormone and corticosteroids can be mentioned, since prolonged use of these substances produces exaggerated protein catabolism.

WARNINGS:

Keep out of reach of children and pets. National Poison Control Center: 0800-333-0160 Buenos Aires, Argentina.

Must be administered by a veterinarian.

Chinfield S.A. is not responsible for problems caused by improper use and/or storage of the product.

STORAGE CONDITIONS:

Store below 30 °C, protected from light.

Do not refrigerate. Do not freeze.

SALES UNDER PRESCRIPTION ONLY

Manufactured and Distributed by: Chinfield S.A. Av. Ader 3205, Munro, Buenos Aires, Argentina

E-Mail: asuntosregulatorios@chinfield.com

www.chinfield.com